

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07H 21/00, C12N 15/11, A61K 31/70		A1	(11) International Publication Number: WO 94/17093
			(43) International Publication Date: 4 August 1994 (04.08.94)
(21) International Application Number: PCT/US94/00902		(74) Agents: McDONNELL, John, J. et al.; Allegetti & Witcoff, Ltd., Ten South Wacker Drive, Chicago, IL 60606 (US).	
(22) International Filing Date: 25 January 1994 (25.01.94)			
(30) Priority Data: 08/009,262 25 January 1993 (25.01.93) US		(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(60) Parent Application or Grant (63) Related by Continuation US 08/009,262 (CON) Filed on 25 January 1993 (25.01.93)		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(71) Applicant (for all designated States except US): HYBRIDON, INC. [US/US]; One Innovation Drive, Worcester, MA 01605 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): KANDIMALLA, Ekambar, R. [IN/US]; Apartment 305, 285 Plantation Street, Worcester, MA 01604 (US). TEMSAMANI, Jamal [MA/US]; Shrewsbury Green Drive 18L, Shrewsbury, MA 01545 (US). AGRAWAL, Sudhir [IN/US]; 61 Lamplighter Drive, Shrewsbury, MA 01545 (US).			
(54) Title: OLIGONUCLEOTIDE ALKYLPHOSPHONATES AND ALKYLPHOSPHONOTHIOATES			
(57) Abstract			
<p>Disclosed is an oligonucleotide analog comprising at least one ribonucleotide alkylphosphonate or alkylphosphonothioate. This analog is preferably from 2 to 60 nucleotides in length and has at least one ribonucleotide substituted at the 2' position of its ribose group. Also disclosed are therapeutic formulations comprising this oligonucleotide analog, methods of inhibiting the expression of a gene from a virus, pathogenic organism, or cell, the expression of which is associated with a disease state, and methods of treating a mammal infected with a virus or pathogenic organism or afflicted with a disorder resulting from the expression of a cellular gene.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

OLIGONUCLEOTIDE ALKYLPHOSPHONATES AND ALKYLPHOSPHONOTHIOATES

FIELD OF THE INVENTION

This invention relates to synthetic oligonucleotides useful for antisense applications. More particularly, this invention relates to synthetic oligonucleotides having
5 modifications in their sugar phosphate backbone which enhance their antisense qualities, and to uses thereof.

BACKGROUND OF THE INVENTION

The sequence specific recognition and binding of various
10 biological molecules to cellular nucleic acid targets is fundamental in molecular biological processes such as gene expression and control. Design of peptide based sequence specific agents for various uses including therapeutic purposes is very difficult as the sequence specific
15 recognition and interaction between proteins and DNA is not very well understood to date. However, it is possible to design and develop sequence-specific oligonucleotides to bind any predetermined sequence on DNA through triple helix formation and on RNA through double helix formation because
20 of the complementary recognition of nucleotide bases via hydrogen bonding interactions. Based on this principle, natural and modified oligonucleotides are used extensively as tools in molecular and cellular biology to determine the role played by specific genes in living systems.

25 In recent years oligonucleotides have been used to block expression of specific genes by interacting with mRNA species in cell culture systems (Uhlmann et al. (1990) *Chem. Rev.* 90:543-584; Helene et al. (1990) *Biochem. Biophys. Acta* 1049:99-125). These "antisense" oligonucleotides can be used as
30 therapeutic agents *in vivo*. For example, antisense oligonucleotides can be used to suppress smooth muscle cell migration or growth in carotid arteries of rats (Simons et al. (1992) *Nature* 359:67-70).

In order to be useful as therapeutic agents, antisense oligonucleotides must be able to be taken up by cells and function therein. Native phosphodiester-linked oligonucleotides are not taken up by cells efficiently because of their polyanionic nature. They are also highly susceptible to nuclease action within the cell (see, e.g., Wickstrom (1986) *Biochem. Biophys. Meth.* 13:97-102), limiting their bioavailability in vivo. To improve cellular uptake and resistance to nuclease action, the sugar phosphate backbone of these oligonucleotides have been modified. For example, oligodeoxyribonucleotides containing methylphosphonate linkages have been prepared by Ts'o et al. (U.S. Patent No. 4,469,863) and Miller et al. (*Biochem.* (1981) 20:1874-1880). Other modifications include phosphorothioates, phosphorodithioates, phosphoramidates, and phosphate esters.

"Chimeric" oligodeoxyribonucleotides have been synthesized which have more than one type of internucleotide linkage, such as those modified linkages described above. For example, Pederson et al. (U.S. Patent No. 5,149,797, the teachings of which are hereby incorporated by reference, disclose chimeric oligonucleotides having an oligonucleotide phosphodiester or oligonucleotide phosphorothioate core sequence flanked by oligonucleotide methylphosphonates or phosphoramidates. Furdon et al. (*Nucleic Acids Res.* (1989) 17:9193-9204) disclose chimeric oligodeoxyribonucleotides having regions of oligonucleotide phosphodiester in addition to either oligodeoxyribonucleotide phosphorothioate or methylphosphonate regions. Quartin et al. (*Nucleic Acids Res.* (1989) 17:(7523-7562) disclose chimeric oligodeoxyribonucleotides having regions of oligonucleotide phosphodiester and oligonucleotide methylphosphonates.

Many of these modified oligonucleotides have contributed to improving the potential efficacy of the antisense

oligonucleotide therapeutic approach. However, certain deficiencies remain that limit the effectiveness of such oligonucleotides as therapeutic agents. For example, Agrawal et al. (*Proc. Natl. Acad. Sci. (USA)* 87:1401-1405) teach that
5 oligodeoxyribonucleotide phosphoramidates when hybridized to RNA do not activate RNase H, the activation of which can be important to the function of antisense oligonucleotides. Also, Agrawal et al. (*Nucleosides & Nucleotides* (1989) 8:5-6) teach that oligodeoxyribonucleotide phosphorothioates have reduced
10 duplex stability when hybridized to RNA.

Thus, there is a need for improved oligonucleotides that overcome the deficiencies of oligonucleotides known in the art. Ideally, such oligonucleotides should be resistant to nucleolytic degradation and should form stable duplexes with
15 RNA.

SUMMARY OF THE INVENTION

In a first aspect, the invention provides synthetic oligonucleotide analogs that have been prepared with non-
20 ionic internucleotide linkages. The subject oligonucleotide analogs are highly nuclease resistant, form a very stable duplex with RNA, and have the requisite hydrophobicity for successful cellular uptake. A requirement of these analogs is that they have at least one ribonucleotide
25 alkylphosphonate or alkylphosphonothioate at any selected position(s) within the oligonucleotide and/or at either or both ends.

For purposes of the invention, the term "oligonucleotide" includes polymers of one or more
30 ribonucleotide monomers connected together or linked to a deoxyribonucleotide monomer by at least one 5' to 3' internucleotide linkage. These linkages include any of the linkages known in the antisense oligonucleotide art. In

addition, the term "oligonucleotide" includes molecules having modified nucleic acid/bases and/or sugars, added substituents, such as diamines, cholesteryl or other lipophilic groups, and 2'-substituted ribonucleoside monomers. The term "oligonucleotide analog" encompasses oligonucleotides with at least one non-phosphodiester internucleotide linkage and/or with modified nucleic acid/bases and/or sugars.

In a preferred embodiment, oligonucleotide analogs according to the invention range from about 2 to about 60 nucleotides in length, and most preferably from about 25 to about 30 nucleotides in length. Thus, oligonucleotides according to the invention will have from 1 to about 59 ribonucleotide alkylphosphonates or alkylphosphonothioates, or a combination of such linkages, and preferably from about 24 to 29 groups.

As used herein, an alkylphosphonate or alkylphosphonothioate is a deoxyribonucleotide or ribonucleotide, or 2'-substituted ribonucleotide, having a phosphonate or phosphonothioate linkage with a 2' alkyl group, including but not limited to, methyl, ethyl, propyl, or butyl group.

In some embodiments of the invention, the oligonucleotide analogs have alkylphosphonate and/or alkylphosphonothioates at some positions within the molecule and natural nucleotide linked by phosphodiester and/or artificial linkages at other positions. These are considered "chimeric" or "hybrid" oligonucleotides.

As used herein, an artificial internucleotide linkage is a non-phosphodiester linkage which links ribonucleotide monomers together from 5' to 3', and which includes, but is not limited to, phosphorothioates, phosphorodithioates, carbonates, phosphate esters, phosphoramidates, and carbamates. In another embodiment of the invention, the

oligonucleotide analogs include at least one, and preferably 6 to 10 2'-substituted ribonucleosides. For purposes of the invention, the term "2'-substituted" means substitution of the 2'-OH of the ribose molecule with, *e.g.*, 2'-O-methyl, 2'-allyl, 2'-aryl, 2'-alkyl, 2'-halo, or 2'-amino, but not with 2'-H, wherein allyl, aryl, or alkyl groups may be unsubstituted or substituted, *e.g.*, with halo, hydroxy, cyano, nitro, acyl, acyloxy, alkoxy, carboxyl, trifluoromethyl, carbalkoxyl, or amino groups.

10 In a second aspect, the invention provides therapeutic pharmaceutical formulations that are effective for treating viral infection, infections by pathogenic organisms, or disease resulting from abnormal gene expression or from the expression of an abnormal gene product. Such therapeutic
15 pharmaceutical formulations include the oligonucleotides according to the first aspect of the invention in a physiologically acceptable carrier.

In a third aspect, the invention provides a method for inhibiting the expression of a viral gene, a gene of a
20 pathogenic organism, or a cellular gene using the oligonucleotide analogs described above. The method includes treating the cells infected with the virus or pathogenic organism in the former two cases, or the cells, generally, in the latter case, with the oligoribonucleotide analog of
25 the invention.

In a fourth aspect, the invention provides a method of treating a mammal afflicted with a disease or disorder resulting from infection with a virus or pathogenic organism, or resulting from the abnormal expression or product of a
30 cellular gene. The method includes administering therapeutic pharmaceutical formulations including the oligonucleotide analogs according to the invention to the mammal in an amount sufficient to form a stable duplex with the nucleic acid of

the virus, pathogen, or gene, thereby inactivating it. Preferable routes of such administration include oral, intranasal, rectal, intravenous, and topical administration. This method of treatment may also include the administration
5 of other therapeutic agents in conjunction with the oligonucleotide analog-containing formulation.

BRIEF DESCRIPTION OF THE DRAWINGS

The foregoing and other objects of the present
10 invention, the various features thereof, as well as the invention itself may be more fully understood from the following description, when read together with the accompanying drawings in which:

15 FIG. 1 is a schematic representation of various phosphate backbone linkages and modifications: FIG. 1A, phosphodiester; FIG. 1B, methylphosphonothioate; FIG. 1C, methylphosphonate; FIG. 1D, phosphorothioate; FIG. 1E, phosphorodithioate; 1F, phosphoramidate; and 1G, phosphate
20 ester;

FIG. 2 is a schematic representation of the synthesis of 3'-phosphoramidites and their subsequent incorporation into an oligonucleotide;
25

FIG. 3 is a graphic representation comparing ion exchange HPLC profiles of 11mer analogs 1 to 3 (SEQ ID NOS:1-3) and control d(A) 11mer;

30 FIG. 4 is a graphic representation of the melting characteristics of duplexes formed from the 25mer analog (SEQ ID NO:4) and complementary strands of (A) DNA and (B) RNA;

FIG. 5 is a graphic representation comparing digestion profiles obtained after snake venom phosphodiesterase digestion of (FIG. 5A) oligonucleotide analog (SEQ ID NO:4) and (FIG. 5B) control oligonucleotide (SEQ ID NO:5);

5

FIG. 6 is an autoradiogram showing the phosphodiesterase I digestion pattern of oligonucleotides 1-3 (SEQ ID NOS:1-3) and d(A)₁₁ (SEQ ID NO:6) at different time intervals; and

10

FIG. 7 is an autoradiogram of a polyacrylamide gel demonstrating the stability of oligonucleotides having SEQ ID NO:4 and SEQ ID NO:5 in the presence of 10% fetal bovine serum.

15

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention provides oligonucleotide analogs characterized by having at least one internucleotide linkage that is an alkylphosphonate or alkylphosphonothioate linkage. The remaining linkages can be phosphodiester linkages or other artificial internucleotide linkages including but not limited to phosphorothioates, phosphorodithioates, phosphoramidates, and phosphate esters. Some of these linkages are depicted in FIG. 1. Such modification of the nucleic acid sugar backbone improves cellular uptake and resistance to nuclease action. The number of alkylphosphonate or alkylphosphonothioate linkages can range from 1 to as many internucleotide linkages as are present in the oligonucleotide. Thus, for purposes of the invention, the term "oligonucleotide alkylphosphonate" or "oligonucleotide alkylphosphonothioate" is intended to encompass every such embodiment.

30

In a preferred embodiment, oligonucleotides according to the invention will range from about 2 to about 60

nucleotides in length, and most preferably from about 25 to about 30 nucleotides in length. Accordingly, in these embodiments, oligonucleotides of the invention will have from 1 to about 59 and from 24 to 29 alkylphosphonate or alkylphosphonothioate internucleotide linkages, respectively.

In some embodiments of the invention, the oligonucleotides have at least one, and preferably 6 to 10 2'-substituted ribonucleoside. Useful 2'-substitutions include, *e.g.*, 2'-O-methyl, 2'-allyl, 2'-aryl, 2'-alkyl, 2'-halo, or 2'-amino, but not with 2'-H, wherein allyl, aryl, or alkyl groups may be unsubstituted or substituted, *e.g.*, with halo, hydroxy, trifluoromethyl, cyano, nitro, acyl, acyloxy, alkoxy, carboxyl, carbalkoxyl or amino groups. These substitutions can be performed by methods well known to the skilled artisan (see, *e.g.*, Goodchild (1990) *Bioconjugate Chem.* 1:165-187). In a particularly useful oligonucleotide, there are 6 or more ribonucleosides and/or 2'-substituted ribonucleosides to enhance duplex stability.

The oligonucleotide alkylphosphonate analogs of the invention can be prepared, for example, by synthesizing 3'-phosphonamidites of 2'-O-Me 5'-DMT ribonucleosides essentially according to the method of Agrawal and Goodchild (*Tetr. Lett.* (1987) 28:3539-3542). This procedure is diagrammed in FIG. 2. The dry phosphonamidites thus produced may then be used on an automated DNA synthesizer to make the required oligonucleotides. Coupling is carried out using suitably protected 2'-O-methyl-ribonucleoside methylphosphonamidites at desired positions. The oligomers so synthesized are cleaved from the controlled pore (CPG) support and then deprotected with ethylenediamine. Representative oligonucleotide analogs of the invention are set forth as SEQ ID NOS:1-4 and 7.

The oligonucleotide methylphosphonothioates of the invention can be prepared by the same methodology used for the synthesis of alkylphosphonates as shown in FIG. 2B except that the oxidant is a 1% solution of 3H-1,2-benzodithiol-3-one 1,1 dioxide (Iyer et al (1990) *J. Org. Chem.* 55:4693-4698) instead of $I_2/H_2O/THF/lutidine$.

The oligonucleotide analogs so produced may then be tested for their ability to hybridize stably with complementary DNA and RNA strands by determining the temperature at which the analog-complementary strand duplexes melted. The melting curves of a representative oligonucleotide methylphosphonate analog which is 25 bases long (a 25mer) duplexed to (A) complementary DNA or to (B) complementary RNA are shown in FIG. 4. The melting temperature (T_m) values obtained with the modified 25mer (SEQ ID NO:4) 4 and the normal 25mer (SEQ ID NO:5) are summarized in TABLE 1. These values are the average of two experiments.

TABLE 1

Thermal Melting Data

Oligomer	Complementary Strand	$T_m, ^\circ C$	% Hyper-Chromicity
25mer analog (SEQ ID NO:4)	DNA	63.7	19.2
	RNA	68.7	15.0
25mer control (SEQ ID NO:5)	DNA	65.7	22.3
	RNA	71.8	18.1

These results suggest that the methylphosphate modification does not hinder the ability of the oligonucleotide to form a duplex with a complementary strand of DNA or RNA, nor does its presence destabilize duplex formation.

To show that the methylphosphonate linkage is incorporated into the oligonucleotide and that the presence of this modification does not prevent subsequent addition and extension of the nucleotide chain, representative oligonucleotide analogs 1-3 (SEQ ID NOS:1-3) with one, two, and three methyl-phosphonate linkages, respectively, were synthesized and studied by ion exchange HPLC. As expected, the unmodified d(A)₁₁ control was eluted at 30 minutes on the Partisil SAX column. Oligonucleotide analogs 1-3 (SEQ ID NO:1-3) were eluted at 23, 20, and 18 minutes, respectively, using a 0 to 50% gradient of solvent A (60% formamide, 40% water, 1 mM potassium dihydrogen phosphonate, pH 6.3) and solvent B (60% formamide, 40% water, 300 mM potassium dihydrogen phosphate, pH 6.3).

The stability of oligonucleotide methylphosphonate analogs was examined in the presence of snake venom phosphodiesterase (SVPD) or phosphodiesterase I, a 3'-exonuclease using the 25mer (SEQ ID NO:4) as the sample. Increase in absorbance due to the hydrolysis of an oligonucleotide was monitored on a spectrometer. The resulting digestion profiles of the 25mer analog (SEQ ID NO:4) and the control 25mer (SEQ ID NO:5) are shown in FIG. 5. 50% of the control 25mer (SEQ ID NO:5) was digested in about 100 seconds. However, the half life of the modified 25mer (SEQ ID NO:4) was measured as 800-900 seconds under the same experimental conditions indicating that the modification is resistant to nuclease action.

The stability of the oligonucleotides was also examined in the presence of phosphodiesterase I. The d(A)₁₁ (SEQ ID NO:6) oligonucleotide was used as the control. The results are shown in FIG. 6. The control (SEQ ID NO:6) was completely degraded into mononucleotides within 5 minutes. The oligonucleotide 1 (SEQ ID NO:1) which contains one

methyolphosphonate linkage, is slightly more stable than d(A)₁₁ (SEQ ID NO:6). However, in 10 minutes, almost all of oligonucleotide 1 (SEQ ID NO:6) was digested. In the case of oligonucleotides 2 and 3 (SEQ ID NOS: 2 and 3) which
5 contain two and three methyolphosphonate linkages, respectively, degradation stopped at nucleotide 6 or 7. This suggests that the phosphodiesterase I may cleave the first methyolphosphonate from the 3'-end but cannot cleave the second one.

10 In another aspect, the invention provides oligonucleotide analogs that are effective in inhibiting viruses, pathogenic organisms, or the expression of cellular genes. The ability to inhibit such agents is clearly important to the treatment of a variety of disease states.
15 Oligonucleotide analogs according to this aspect of the invention share the characteristics of the above-described oligonucleotide analogs, and also have a nucleotide sequence that is complementary to a nucleic acid sequence that is from a virus, a pathogenic organism, or a cellular gene.
20 Preferably such oligonucleotides are from about 6 to about 50 nucleotides in length. For purposes of the invention, the term "oligonucleotide sequence that is complementary to a nucleic acid sequence" is intended to mean an oligonucleotide sequence (2 to about 50 ribonucleotides) that hybridizes to
25 the nucleic acid sequence under physiological conditions, *e.g.*, by Watson-Crick base pairing (interaction between oligonucleotide and single-stranded nucleic acid), by Hoogsteen base pairing (interaction between oligonucleotide and double-stranded nucleic acid), or by any other means.
30 Such hybridization under physiological conditions is measured as a practical matter by observing interference with the function of the nucleic acid sequence.

The nucleic acid sequence to which an oligoribonucleotide analog of the invention is complementary will vary, depending upon the agent to be inhibited or the nucleic acid to be targeted. In many cases the nucleic acid sequence will be a viral nucleic acid sequence. The use of antisense oligonucleotides to inhibit various viruses is well known, and has recently been reviewed in Agrawal (*Tibtech* (1992) 10:152-15). Viral nucleic acid sequences that are complementary to effective antisense oligonucleotides have been described for many viruses, including human immunodeficiency virus type 1 (U.S. Patent No. 4,806,463, the teachings of which are herein incorporated by reference), *Herpes simplex* virus (U.S. Patent No. 4,689,320, the teachings of which are hereby incorporated by reference), Influenza virus (U.S. Patent No. 5,194,428, the teachings of which are hereby incorporated by reference), and Human papilloma virus (Storey et al. (1991) *Nucleic Acids Res.* 19:4109-4114). Oligonucleotides according to the invention can have sequences complementary to any of these viral nucleic acid sequences. Additional viruses that have known nucleic acid sequences against which antisense oligonucleotides can be prepared include Foot and Mouth Disease Virus (see Robertson et al. (1985) *J. Virol.* 54:651; Harris et al. (1980) *J. Virol.* 36:659); Yellow Fever Virus (see Rice et al. (1985) *Science* 229:726), Varicella-Zoster Virus (see Davison and Scott (1986) *J. Gen. Virol.* 67:2279); and Cucumber Mosaic Virus (see Richards et al. (1978) *Virol.* 89:395).

Alternatively, oligonucleotides according to the invention can have a nucleic acid sequence complementary to a nucleic acid sequence of a pathogenic organism. The nucleic acid sequences of many pathogenic prokaryotes have been described, and include the malaria organism, *Plasmodium falciparum*, and many other pathogenic bacteria. Examples of

pathogenic eucaryotes having known nucleic acid sequences against which antisense oligonucleotides can be prepared include *Trypanosoma brucei gambiense* and *Leishmania* (see Campbell et al. (1984) *Nature* 311:350); *Fasciola hepatica* (see Zurita et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:2340). Antifungal oligonucleotides can be prepared using a target hybridizing region having an oligonucleotide sequence that is complementary to a nucleic acid sequence from, e.g., the chitin synthetase gene. Likewise, antibacterial oligonucleotides can be prepared using, e.g., the alanine racemase gene.

The oligonucleotides according to the invention alternatively can have an oligonucleotide sequence complementary to a cellular gene or gene transcript, the abnormal expression or product of which results in a disease state or disorder. The nucleic acid sequences of several such cellular genes have been described, including the prion protein (Stahl et al. (1991) *FASEB J.* 5:2799-2807), the amyloid-like protein associated with Alzheimer's disease (U.S. Patent No. 5,015,570, the teachings of which are hereby incorporated by reference), and various well-known oncogenes and proto-oncogenes, such as *c-myb*, *c-myc*, *c-abl*, and *n-ras*. In addition, oligonucleotides that inhibit the synthesis of structural proteins or enzymes involved largely or exclusively in spermatogenesis, sperm motility, the binding of the sperm to the egg or any other step affecting sperm viability may be used as contraceptives for men. Similarly, contraceptives for women may be oligoribonucleotides that inhibit the synthesis or function of proteins or enzymes involved in ovulation, fertilization, implantation, or in the biosynthesis of hormones involved in those processes.

The oligonucleotide analogs of the invention also can be used to control hypertension. Such analogs suppress the synthesis of angiotensin converting enzyme or related enzymes

in the renin/angiotensin system. Similarly, platelet aggregation can be controlled by suppression of the synthesis of enzymes necessary for the synthesis of thromboxane A₂ for use in myocardial and cerebral circulatory disorders, infarcts, arteriosclerosis, embolism and thrombosis. Deposition of cholesterol in arterial wall can be inhibited by suppression of the synthesis of fattyacyl co-enzyme A:cholesterol acyl transferase in arteriosclerosis. Inhibition of the synthesis of cholinephosphotransferase may be useful in hypolipidemia.

There are numerous neural disorders in which hybridization arrest can be used to reduce or eliminate adverse effects of the disorder. For example, suppression of the synthesis of monoamine oxidase can be used in Parkinson's disease; suppression of catechol-o-methyl transferase can be used to treat depression; and suppression of indole N-methyl transferase can be used in treating schizophrenia.

Suppression of selected enzymes in the arachidonic acid cascade which leads to prostaglandins and leukotrienes may be useful in the control of platelet aggregation, allergy, inflammation, pain and asthma.

Suppression of the protein expressed by the multidrug resistance (*mdr*) gene, which is responsible for development of resistance to a variety of anti-cancer drugs and is a major impediment in chemotherapy may prove to be beneficial in the treatment of cancer.

Oligonucleotide sequences complementary to nucleic acid sequences from any of these genes can be used for oligonucleotides according to the invention, as can be sequences complementary to any other cellular gene or gene transcript, the abnormal expression or product of which results in a disease state or disorder.

Antisense regulation of gene expression in plant cells has been described in U.S. Patent No. 5,107,065, the teachings of which are hereby incorporated by reference.

5 In another aspect, the invention provides a method for inhibiting the gene expression of a viral gene, a gene of a pathogenic organism, or a cellular gene using the oligonucleotide analogs described above. The method includes treating the cells infected with the virus or pathogenic
10 organism in the former two cases, or the cells generally in the latter case, with the analogs of the invention in an amount sufficient to hybridize with available copies of the nucleic acid to which it is complementary. Treatment may be accomplished by directly administering the analogs to the cells, which may be cultured or in an organism, as described
15 below.

In yet another aspect, the invention includes therapeutic pharmaceutical formulations that are effective for treating virus infection, infection by pathogenic organisms, or a disorder or disease resulting from abnormal
20 gene expression or from the expression of an abnormal gene product. Such therapeutic pharmaceutical formulations contain the oligonucleotides according to the first aspect of the invention in a physiologically acceptable carrier.

As used herein, a "physiologically acceptable carrier"
25 includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or
30 agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

These pharmaceutical formulations can be used to practice yet another embodiment of the invention, namely, a method of treating a mammal having a disease or disorder resulting from infection with a virus or pathogenic organism, or from the abnormal expression or product of a cellular gene. The method comprises administering the therapeutic pharmaceutical formulation of the invention to the afflicted mammal in an amount sufficient to enable the binding of the oligonucleotide to the complementary nucleic acid to which it has been targeted. In this way, the oligonucleotide inhibits the expression and replication of the gene from the virus, pathogenic organism, or expression of the abnormal cellular gene.

Effective dosages of the oligonucleotide and modes of its administration in the treatment of various disorders such as AIDS can be determined by routine experimentation. The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile. It must be stable under the conditions of manufacture and storage and may be preserved against the contaminating action of microorganisms, such as bacterial and fungi. The carrier can be a solvent or dispersion medium. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents. Prolonged absorption of the injectable therapeutic agents can be brought about by the use of the compositions of agents delaying absorption.

The oligonucleotide in the carrier may be administered by any useful route including intravenous or intraperitoneal injection, or by intranasal, rectal, oral, transdermal, or subcutaneous administration. Sterile injectable solutions

are prepared by incorporating the oligonucleotide in the required amount in the appropriate solvent, followed by filtered sterilization.

5 The method of the invention may be used to treat variety of viral diseases including AIDS, ARC, oral or genital herpes, papilloma warts, flu, foot and mouth disease, yellow fever, chicken pox, shingles, HTLV-leukemia, and hepatitis. Among fungal diseases treatable by the method are candidiasis, histoplasmosis, cryptococcocis, blastomycosis, 10 aspergillosis, sporotrichosis, chromomycosis, dermatophytosis and coccidioidomycosis. The method can also be used to treat rickettsial diseases (e.g., typhus, Rocky Mountain spotted fever), as well as sexually transmitted diseases caused by *Chlamydia trachomatis* or *Lymphogranuloma venereum*. A variety of 15 parasitic diseases can be treated by the method, including amebiasis, Chagas' disease, toxoplasmosis, pneumocystosis, giardiasis, cryptosporidiosis, trichomoniasis, and *Pneumocystis carini* pneumonia, worm (helminthic diseases) such as ascariasis, filariasis, trichinosis, schistosomiasis, and 20 nematode or cestode infections. Malaria can be treated by the method regardless of whether it is caused by *P. falciparum*, *P. vivax*, *P. orale*, or *P. malariae*.

The infectious diseases identified above can all be treated by the method of the invention because the infectious 25 agents for these diseases are known, and thus oligonucleotides according to the invention can be prepared to have a nucleic acid sequence that is complementary to a nucleic acid sequence essential for the propagation of the infectious agent (e.g., an essential gene).

30 Other disease states or conditions that are treatable by the method result from an abnormal expression or product of a cellular gene. These conditions have been discussed above.

The following example illustrates the preferred mode of making and practicing the present invention, but is not meant to limit the scope of the invention since alternative methods may be utilized to obtain similar results.

5

EXAMPLES

1. Synthesis of Ribonucleoside Phosphonamidites

3'-phosphonamidites of 2'-O-methyl 5'-DMT ribonucleosides were synthesized as described by Agrawal and Goodchild (Agrawal et al. (1987) *Tetr. Lett.* 28:3539-3542). This procedure is shown in FIG. 2. Briefly, to 0.024 mmol of 2'-O-methyl 5'-DMT ribonucleoside is added 0.4 mmol of Hunig's base (ethyl diisopropyl amine) in 5 ml of anhydrous dichloromethane. This is cooled in an ice bath for 10 minutes. 0.4 mmol of methyl N, N-diisopropylchloro phosphoramidite is added slowly in 2 ml of dichloromethane. The temperature of the solution is allowed to rise to ambient level, after which time it is left at room temperature for 30 minutes. The reaction is then quenched by adding 30 ml of ethyl acetate. The mixture is extracted with saturated sodium bicarbonate and brine solutions. The organic layer is dried over anhydrous sodium sulfate and evaporated to dryness. The resulting phosphonamidites are dissolved in a small amount of ethyl acetate and then precipitated by adding the ethyl acetate solution to a large volume of pentane at -30°C. The precipitate is collected by centrifugation and dried.

2. Oligonucleotide Synthesis

Dry phosphoramidites are dissolved in DNA synthesis grade acetonitrile at a concentration of 30 mg/ml and are used on an automated DNA synthesizer to make the required oligonucleotides. The oligonucleotide synthesis is carried

out on Milligen/Bioscience 8700 automatic synthesizer using RNA synthesis cycle at 1 μ M scale using the phosphoramidite approach (Agrawal et al. (1987) *Tetra. Lett.* 28:3539-3542). Coupling is carried out using suitably protected 2'-O-methyl-
5 ribonucleoside methylphosphonamidites at required positions for oligonucleotide analogs 1-4 (SEQ ID NOS:1-4).

3. Deprotection

The oligomers are cleaved from CPG using 1:1 ammonium
10 hydroxide-ethanol mixture for 1 hour at room temperature and then deprotected with ethylenediamine at room temperature for 6 hours.

4. Purification and Assay

15 The 5'-DMT protected oligomers are purified by reverse phase chromatography using reverse phase (RP) C18 HPLC column using 0.1M ammonium acetate and a 1:4 ratio of 0.1M ammonium acetate-acetonitrile gradient from 0 to 50%. The oligomers are detritylated with 80% acetic acid at room temperature for
20 1 hour and then desalted on a C₁₈ Sep-pack cartridge (Agrawal et al. (1987) *Tetra. Lett.* 28:3539-3542). The oligomers are then purified on 20% polyacrylamide denaturing gel. The oligomers are again desalted on sep-pack cartridge and used for further studies.

25 To show that the methylphosphonate linkage is incorporated into the oligonucleotide and that the presence of this modification does not prevent subsequent addition and extension of the nucleotide chain, analogs 1-3 (SEQ ID NOS:1-3) were assayed by ion exchange HPLC. A 0 to 50% gradient
30 of solvents A (60% formamide, 40% water, 1 mM potassium dihydrogen phosphate, pH 6.3) and B (60% formamide, 40% water, 300 mM potassium dihydrogen phosphate, pH 6.3) was used.

5. Hybridization Studies

Hybridization properties of the modified 25mer (SEQ ID NO:4) were studied by performing thermal melting experiments with DNA and RNA complementary strands. Typically, the modified oligonucleotide and the complementary strand (1:1) were dried in an Eppendorf tube and dissolved in 1 ml of 100 mM NaCl and 10 mM phosphate buffer, pH 7.4. The solution is heated to 80°C for 5 minutes and allowed to come to room temperature slowly. The T_m is measured on Perkin Elmer Lambda 2 UV/Vis Spectrometer interfaced with a Digital DECstation 316sx computer. The resulting melting curves of the 25mer analog with a complementary strand of A(DNA) and B(RNA) are shown in FIG. 4. The melting temperature (T_m) values obtained with the modified 25mer (SEQ ID NO:4) and the normal control 25mer (SEQ ID NO:5) are summarized in TABLE I. These values are the average of two experiments.

6. Stability of Oligonucleotide Analogs in Nucleases

The stability of the modified 25mer (SEQ ID NO:4) was examined in the presence of snake venom phosphodiesterase (SVPD), or phosphodiesterase I, a 3'-exonuclease. Increase in absorbance due to the hydrolysis of an oligonucleotide was monitored on a Perkin Elmer Lambda 2 UV/Vis Spectrometer. In a typical experiment, 0.2 A_{260} units of modified oligonucleotide was dissolved in 0.5 ml of 10 mM $MgCl_2$, 10 mM Tris buffer, pH 8.3, and equilibrated at 37°C for 5 minutes. 1.5×10^{-3} units of SVPD was added, and the rate of digestion followed by recording change in absorbance at 260 nm with time. The resulting digestion profiles are shown in FIG. 5.

The stability of the oligonucleotides 1-3 (SEQ ID NOS:1-3) with reference to the control normal oligonucleotide d(A)₁₁ (SEQ ID NO:6) was studied with 3'-phosphodiesterase I by polyacrylamide gel electrophoresis. The oligoribonucleotides

(about 100 ng) were labelled at their 5'end with ^{32}P using γ - ^{32}P -ATP and T4 polynucleotide kinase (Sambrook et al., *Molecular Cloning, A Laboratory Manual* (2d ed.) Cold Spring Harbor Laboratory Press, 1989). Each 5'-end labelled oligoribonucleotide (3
5 μg) was treated with 1 μl of phosphodiesterase I (1.5×10^{-4} Units) in 30 μl of buffer containing 50 mM Tris-HCl, pH 7.5, 5 mM MgCl_2 , 5 mM DTT, 50 $\mu\text{g/ml}$ BSA. 4 μl aliquot of sample was removed from the reaction mixture at 0, 2, 5, and 10
10 minutes intervals. The aliquots were heated at 90°C for 5 minutes and separated by electrophoresis on a 20% polyacrylamide-8M urea gel. The autoradiogram developed is shown in FIG. 6.

The stability of the modified 25mer analog (SEQ ID NO:4) with reference to the control 25mer oligonucleotide (SEQ ID
15 NO:5) was studied in cell culture media containing 10% fetal bovine serum. The oligonucleotides (about 100 ng) were labelled at the 5'-end with ^{32}P as described in the preceding paragraph. About 10 ng of each of the labelled oligonucleotides (specific activity 10^6 dpm/10 ng), were
20 separately incubated in 300 μl of SMEM cell culture media (GIBCO) containing 10% fetal bovine serum. At each time point (0, 1, 2, and 4 hr), 50 μl aliquots were removed, treated with 10 μl of proteinase K, phenol-chloroform extracted, and then precipitated with ethanol. The resulting
25 reaction mixtures were analyzed by electrophoresis on a 20% polyacrylamide gel containing 8M urea. The developed autoradiogram is shown in FIG. 7.

The results show that the modified 25mer oligonucleotide analog is intact for 4 hours in considerable amounts compared
30 to the control 25mer which is degraded into shorter sequences in less than 1 hour.

7. Inhibition of HIV

Oligonucleotide methylphosphonates are tested for their ability to inhibit HIV-1 in tissue culture as follows. H9 lymphocytes are infected with HIV-1 virions ($\approx 0.01 - 0.1$ TCID₅₀/cell) for one hour at 37°C. After one hour, unadsorbed virions are washed and the infected cells are divided among wells of 24-well plates. To the infected cells, an appropriate concentration (from stock solution) of oligonucleotide analog is added to obtain the required concentration in 2 ml medium. The cells are then cultured for four days. At the end of four days, level of HIV-1 expression is measured by synthetic formation, p24 expression, and reverse transcriptase activity. The level of expression of p24 in infected cells treated with oligonucleotide analog is compared with the level detected in untreated infected cells.

Those skilled in the art will recognize, or be able to ascertain, using no more than routine experimentation, numerous equivalents to the specific substances and procedures described herein. Such equivalents are considered to be within the scope of this invention, and are covered by the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANTS: HYBRIDON, INC.
- (ii) TITLE OF INVENTION: Oligonucleotide
Alkylphosphonates and
Alkylphosphonothioates
- 10 (iii) NUMBER OF SEQUENCES: 7
- (iv) CORRESPONDENCE ADDRESS:
15 (A) ADDRESSEE: Allegretti & Witcoff, Ltd.
(B) STREET: 10 S. Wacker Drive, Suite 3000
(C) CITY: Chicago
(D) STATE: Illinois
(E) COUNTRY: U.S.A.
(F) ZIP: 60606
- 20 (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
25 (D) SOFTWARE: PatentIn Release #1.0,
Version #1.25
- (vi) CURRENT APPLICATION DATA:
30 (A) APPLICATION NUMBER: US/08/009,262
(B) FILING DATE: 01/25/93
(C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
35 (A) NAME: Greenfield, Michael S.
(B) REGISTRATION NUMBER: 37,142
(C) REFERENCE/DOCKET NUMBER: 92,774-A
- (ix) TELECOMMUNICATION INFORMATION:
40 (A) TELEPHONE: 312/715-1000
(B) TELEFAX: 312/715-1234

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 11 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

10

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

15

(ix) FEATURE:

- (A) NAME/KEY:
(B) LOCATION:
(D) OTHER INFORMATION: /product=
20 "phosphodiester linkages between bases
6-7; U is 2-O-methyl"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

25 TTTTUTTTT T 11

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- 30 (A) LENGTH: 11 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

40

(ix) FEATURE:

- (A) NAME/KEY:
(B) LOCATION:
45 (D) OTHER INFORMATION: /product=
"phosphodiester linkages between
nucleosides 3-4, 9-10; U is 2-O-
methyl"

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TTUTTTTTUT T

11

5

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 11 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

15

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

20

(ix) FEATURE:

- (A) NAME/KEY:
(B) LOCATION:
(D) OTHER INFORMATION: /product=
"phosphodiester linkages between
25 nucleosides 3-4, 6-7, 9-10; U is 2-O-
methyl"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

30

TTUTTUTTUT T

11

(2) INFORMATION FOR SEQ ID NO:4:

35

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: mRNA

(iii) HYPOTHETICAL: NO

45

(iv) ANTI-SENSE: YES

(ix) FEATURE:

- (A) NAME/KEY:
- (B) LOCATION:
- (D) OTHER INFORMATION: /product=
"phosphodiester linkages between
nucleosides 21-22, 22-23, 23-24, and
24-25; bases 21-24 are 2-O-methyl"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CTCTCGCACC CATCTCTCTC CUUCT

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CTCTCGCACC CATCTCTCTC CTTCT

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6

AAAAAAAAA A

(2) INFORMATION FOR SEQ ID NO:7:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 10 (ii) MOLECULE TYPE: mRNA
(iii) HYPOTHETICAL: NO
(iv) ANTI-SENSE: YES
- 15 (ix) FEATURE:
(A) NAME/KEY:
(B) LOCATION:
(D) OTHER INFORMATION: /product=
20 "phosphodiester linkages between
nucleosides 7-8; C and U are
2-0-methyl"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:
25 CCUUCCT 8

What is claimed is:

- 5 1. An oligonucleotide analog comprising at least one ribonucleotide alkylphosphonate or alkylphosphonothioate.
2. The oligonucleotide analog of claim 1 having from 2 to 60 nucleotides.
- 10 3. The oligonucleotide analog of claim 2 having from about 25 to about 30 nucleotides.
- 15 4. The oligonucleotide analog of claim 2 having from about 2 to 60 alkylphosphonates or alkylphosphonothioates.
- 20 5. The oligonucleotide analog of claim 4 having from about 25 to 30 alkylphosphonates or alkylphosphonothioates.
- 25 6. The oligonucleotide analog of claim 2 having from about 7 to 10 alkylphosphonothioates.
- 30 7. The oligonucleotide analog of claim 1 wherein at least one ribonucleotide is substituted at the 2' position of the ribose group with one of the group consisting of 2'-O-methyl, 2'-allyl, 2'-aryl, 2'-alkyl, 2'-halo, and 2'-amino.
8. The oligonucleotide analog of claim 1 wherein at least one nucleotide is a 2'-substituted ribonucleotide.

9. The oligonucleotide analog of claim 8 wherein the alkylphosphonate or alkylphosphonothioate is selected from the group consisting of methyl, ethyl, propyl, butyl, and alkylphosphonate or alkylphosphonothioate.

10. The oligonucleotide analog of claim 9 wherein the 2'-substituted ribonucleotide is further substituted at its 2'-allyl, 2'-aryl, or 2'-alkyl with one of the group consisting of halo, hydroxy, cyano, nitro, acyl, acyloxy, alkoxy, carboxyl, trifluoromethyl, carbalkoxyl, and amino groups.

11. The oligonucleotide analog of claim 1 comprising a nucleotide sequence that is complementary to a nucleic acid sequence from a virus, pathogenic organism, or cellular gene or cellular gene transcript, the abnormal expression of which results in a disease state.

12. A therapeutic pharmaceutical formulation comprising:

- (a) the oligonucleotide of claim 1; and
- (b) a physiologically acceptable carrier.

13. A method of inhibiting the expression of a gene from a virus, pathogenic organism, or cell, the expression of which is associated with a disease state, the method comprising the steps of:

- (a) providing the oligonucleotide analog of claim 11; and
- (b) treating the virus, pathogenic organism or cell with the oligoribonucleotide analog,

the oligonucleotide analog hybridizing with the or mRNA transcribed from the gene, thereby inhibiting expression of the gene.

5 14. A method of treating a mammal infected with a virus or pathogenic organism or afflicted with a disorder resulting from the expression of a cellular gene, the method comprising the steps of:

10 (a) providing the therapeutic formulation of claim 12; and
(b) administering the therapeutic formulation to the mammal in an amount sufficient to allow the oligonucleotide analog to hybridize to a complementary nucleic acid sequence of the
15 gene, thereby inhibiting the expression of the gene.

20 15. The method of claim 14 wherein the administering step (b) comprises administering the therapeutic formation by a route selected from the group consisting of intravenous or intraperitoneal injection, and intranasal, rectal, oral, transdermal, or subcutaneous administration.

25 16. The method of claim 14 wherein the administering step (b) further comprises administering other therapeutic agents to the mammal.

FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D




FIG. 1E




FIG. 1F






FIG. 1G





- X = O when oxidant is $\text{I}_2/\text{H}_2\text{O}/\text{THF}/\text{Lutidine}$;
- X = S when oxidant is 3H-1,2-benzodithiol-3-one 1,1-dioxide

FIG. 2




FIG. 3




FIG. 4

FIG. 5A




FIG. 5B





FIG. 6

**FIG. 7**

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 94/00902

A. CLASSIFICATION OF SUBJECT MATTER
IPC 5 C07H21/00 C12N15/11 A61K31/70

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 5 C07H C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US,A,4 469 863 (TS'O ET AL.) 4 September 1984 see figures 8,9 see column 4, line 17 - line 44 ---	1-5,7-16
X	EP,A,0 386 987 (GEN-PROBE INCORPORATED) 12 September 1990 see the whole document, but especially, examples 1,2,7,9-12 & WO,A,90 11322 (GEN-PROBE INCORPORATED) ---	1,2,4,6, 11-16
X	WO,A,92 04358 (GEN-PROBE INCORPORATED) 19 March 1992 see the whole document ---	1,2,4,6, 11-16
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

28 June 1994

Date of mailing of the international search report

01.07.94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 PatentAan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Scott, J

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 94/00902

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 136 543 (HOECHST AKTIENGESELLSCHAFT) 10 April 1985 see page 1, line 1 - line 9 see claim 1 see table 6 ---	1,2
X	WO,A,91 06556 (GILEAD SCIENCES, INC.) 16 May 1991 see the whole document ---	1-5,7-10
X	TETRAHEDRON LETTERS. vol. 32, no. 37 , 9 September 1991 , OXFORD GB pages 4981 - 4984 J.HELINSKI ET AL. 'N,N-Diisopropyl-O-p-nit rophenyl-P-methylphosphonoamidate: Novel Difunctional Piii Reagent in Oligonucleoside Methylphosphonate Synthesis Containing 4-Nitrophenoxy Group.' see page 4984, line 17 - line 24 ---	1,2
X	TETRAHEDRON LETTERS. vol. 33, no. 17 , 21 April 1992 , OXFORD GB pages 2357 - 2360 H.C.P.F.ROELEN ET AL. 'Synthesis of Alkylphosphon(othio)ate Analogues of DNA.' see the whole document ---	1-3,11
A	WO,A,92 20967 (HYBRIDON, INC.) 26 November 1992 ---	1,11
A	WO,A,91 16902 (HYBRIDON, INC.) 14 November 1991 see the whole document ---	1,11-16
P,X	EP,A,0 552 766 (HOECHST AKTIENGESELLSCHAFT) 28 July 1993 see the whole document ---	1-16
E	WO,A,94 02499 (HYBRIDON, INC.) 3 February 1994 see the whole document -----	1-7, 11-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 94/00902

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4469863	04-09-84	US-A- 4511713 US-A- 4757055	16-04-85 12-07-88
EP-A-0386987	12-09-90	CA-A- 2011430 JP-T- 4503956 WO-A- 9011322 US-A- 5252723	06-09-90 16-07-92 04-10-90 12-10-93
WO-A-9011322	04-10-90	CA-A- 2011430 EP-A- 0386987 JP-T- 4503956 US-A- 5252723	06-09-90 12-09-90 16-07-92 12-10-93
WO-A-9204358	19-03-92	AU-B- 642673	28-10-93
EP-A-0136543	10-04-85	DE-A- 3332068 AU-B- 601257 AU-A- 1138488 AU-B- 570266 AU-A- 3275684 CA-A- 1235079 DE-A- 3475307 JP-B- 5031560 JP-A- 60072899 US-A- 5166330	21-03-85 06-09-90 14-07-88 10-03-88 14-03-85 12-04-88 29-12-88 12-05-93 24-04-85 24-11-92
WO-A-9106556	16-05-91	AU-A- 6715790 EP-A- 0497875 JP-T- 5504552	31-05-91 12-08-92 15-07-93
WO-A-9220967	26-11-92	NONE	
WO-A-9116902	14-11-91	US-A- 5194428 EP-A- 0527906	16-03-93 24-02-93
EP-A-0552766	28-07-93	AU-A- 3191093 CA-A- 2087818	29-07-93 23-07-93
WO-A-9402499	03-02-94	AU-B- 4688393	14-02-94